



# SALMONELLA ENTERICA SPECIES ISOLATED FROM LOCAL FOODSTUFF AND PATIENTS SUFFERING FROM FOODBORNE ILLNESS: SURVEILLANCE, ANTIMICROBIAL RESISTANCE AND MOLECULAR DETECTION

Zahraa A. AlShaheeb<sup>1</sup>, Zaid A. Thabit<sup>2\*</sup>, Asma G. Oraibi<sup>3</sup>, Ahmed A. Baioumy<sup>4</sup>, Tarek Gamal Abdelmaksoud<sup>4</sup>

<sup>1</sup> Iraqi Ministry of Health, Baghdad, Iraq

<sup>2</sup> Environmental Biotech Department, Biotechnology Research Center, Al-Nahrain University, Baghdad, Iraq

<sup>3</sup> Plant Biotechnology Department, College of Biotechnology, Al-Nahrain University, Baghdad, Iraq

<sup>4</sup> Food Science Department, Faculty of Agriculture, Cairo University, Giza, Egypt

**Keywords:** foodborne pathogen, frozen chicken meat, azithromycin resistance gene *mphA*.

## Abstract

The aim of this study was to determine the prevalence of *Salmonella enterica* in raw chicken meat, eggs, and ready-to-eat foods containing poultry products and among patients suffering from diarrhea as a result of ingestion of this foodborne pathogen in Baghdad, Iraq. It assesses the antibiotics susceptibility, virulence and pathogenicity of *S. enterica* isolates. Thirteen *Salmonella* spp. isolates from foodstuff and seven from clinical patients were recovered from 80 and 20 samples, respectively. Isolates from foodstuff samples displayed the highest resistance to nalidixic acid (69.23%), followed by chloramphenicol (53.84%). *Salmonella* spp. isolated from clinical samples showed resistance to both azithromycin and cefotaxime at the same percentage level (71.42%). The results of antibiotic resistance gene amplification (*gyrA*, *mphA*) were analyzed and showed that these genes were present in 100% and 50% of phenotypically resistant isolates, respectively. Virulence genes *invA*, *avrA*, and *sipB* were found on average in 86% of food isolates, accounting for 69.2%, 92.3%, and 95%, respectively. In addition, the detection of these virulence genes among clinical isolates showed their presence at the same level (85.7%). Our study revealed that unhygienic chicken slaughterhouses and lack of food safety management are strong indicators of a high probability of the *Salmonella* presence in our food products in the Iraqi markets.

**For citation:** AlShaheeb, Z., A., Thabit, Z.A., Oraibi, A.G., Baioumy, A.A., Abdelmaksoud, T.G. (2023). *Salmonella enterica* species isolated from local foodstuff and patients suffering from foodborne illness: Surveillance, antimicrobial resistance and molecular detection. *Theory and Practice of Meat Processing*, 8(2), 112-123. <https://doi.org/10.21323/2414-438X-2023-8-2-112-123>

## Introduction

The World Health Organization (WHO) has stated that foodborne diseases remain a significant problem and can show severity among children, the elderly, and people with immunosuppression [1]. More than 250 different foodborne illnesses are caused by various microbial pathogens and toxins [2], poisonous chemicals, or bio-toxins. *Salmonella* infections are a critical epidemiological and economic problem worldwide [4,5]. For example, the European Food Safety Authority (EFSA) reports that *Salmonella* spp. is recognized to be the second most common cause of human diseases and food poisoning associated with contaminated food [6]. *Salmonella* is a genus of gram-negative bacteria that belongs to the *Enterobacteriaceae* family and can infect various animal hosts [7]. Moreover, *Salmonella* can survive under harsh conditions for about a year in frozen meat [8]. *Salmonella* exists in different environments such as soil, water systems, sewage, and the gut flora of several animals [9]. The members of the genus *Salmonella* are categorized on the basis epidemiology, host range, biochemical reactions, and structures of the O, H, and Vi antigens [10]. *Salmonella*, according to their DNA relatedness, can be classified into two species; *S. bongori*, which populates

cold-blooded animals, and *S. enterica*, which is able to inhabit both cold and warm-blooded hosts [11]. Strains belonging to *S. enterica* subsp. *enterica* are responsible for almost 99% of *Salmonella* infections in people and warm-blooded animals [12].

Antimicrobial drugs are utilized to prevent microbial infectious diseases in both humans and animals [7] as well as in animal feed for prophylaxis, therapeutics, and growth promotion [13]. The antimicrobial resistance of *Salmonella*, particularly multidrug resistance (MDR), has become a significant worldwide problem [14]. For example, more recently, *S. enterica* in food-producing animals has shown high MDR resulting in a global problem due to the widespread use of antibiotic drugs [13]. The excessive use of the same antibiotic drugs as a treatment in clinical and veterinary medicine may lead to emergence of resistant strains that can easily be transmitted to the human population from animal products, which is a serious public health problem/concern [15–17]. In Iraq, there is currently a widespread lack of food safety aspects including domestic production of poultry and its products, as well as restaurants and this is leading to the increased risk of exposure to *Salmonella* infection. In Baghdad and central

Iraq, non-typhoidal *Salmonella* was indicated as the second cause of gastroenteritis in children after enteric viruses [18]. Non-typhoidal *Salmonella* was recovered from 10.3% of diarrheal stool samples from children under the age of 5 years in a recent study in southern Iraq [18]. To our knowledge, there have been no published studies on the molecular epidemiology of non-typhoidal *Salmonella* isolated from local hens in Iraq. Given the importance of this pathogen to worldwide health, the current study presents an evaluation of the molecular detection of virulence and antibiotic resistance genes of *Salmonella enterica* from Iraqi foodstuff. We focused on three virulence genes of the SPI-1 region involved in *Salmonella* pathogenicity (*avrA*, *invA*, and *sipB*), which play a critical role in the initial invasion of the host organism and cause salmonellosis infection as previously mentioned in [19–21]. Therefore, the aim of this study was to determine the prevalence of *Salmonella enterica* in raw chicken meat, eggs, and ready-to-eat foods containing poultry products and in patients suffering from diarrhea as a result of ingestion of this foodborne pathogen in Baghdad (Iraq). It assesses the antibiotic susceptibility as well as virulence and pathogenicity.

**Materials and methods**

*Sample collection*

From October 2020 to May 2021, one hundred samples were collected in Baghdad, including 80 samples of Iraqi raw and ready-to-eat food from eight processing points (frozen whole 9-piece chicken=10; frozen chicken breasts=10; frozen chicken thighs=10; eggshell=10; cooked chicken shawarma=10; cooked chicken tika=10; ready-to-eat cake cream=10; food appetizers containing chicken derivatives=10) from local supermarkets, and 20 samples from clinically suspected patients with foodborne diseases from private medical laboratories.

*Salmonella isolation*

Bacteria were isolated according to the ISO 6579–1:2017(E) procedure [22]. An analytical unit of 25 grams from each food sample chopped finely and fecal samples from patients taken with sterilized cotton swabs were inoculated into sterile flasks with 225mL of Buffered Peptone Water (BPW) broth (Himedia, India). The flasks were incubated at 37 °C for 18 hours. An amount of 0.1mL from pre-enriched culture was transferred to 10ml of Rappaport

Vassiliadis (RV) broth medium (Himedia, India), thoroughly mixed and incubated at 41±0.5 °C for 24 hours. A loopful from the incubated selective enrichment broth culture was streaked on Xylose Lysine Deoxycholate (XLD) agar plate and *Salmonella-Shigella* (SS) agar (Himedia, India). Both agar plates were incubated at 37 °C for 24 hours.

*Salmonella identification*

Suspected colonies with *Salmonella* morphology from each plate were identified biochemically using the VITEK-2 system (bioMérieux, France). In addition, isolates were investigated by the conventional method in the Iraqi National Centre for *Salmonella* at the Central Public Health Laboratories in Baghdad using the serological test (Anti-*Salmonella* H test) (Sifin/ Germany) kits that were designed for the use in examining the H-antigens of *Salmonella* strains via slide agglutination in Baghdad’s Central Public Health Laboratory (CPHL). Bacteria from 16–20-hour-old subculture (nutrient) agar were applied to a clean microscope slide and mixed well with a drop of 25µl of anti-*Salmonella* H reagent (test serum), then slowly stirred with a sanitized stick. The slide was put on a dark surface and visible agglutination was observed in the case of the positive reaction, while a negative result was seen as a homogeneous milky turbid suspension. On the same slide, the positive and negative controls were tested in the same way. Typical *Salmonella* phenotypes were further confirmed by single-step PCR for the 16S rRNA gene of *S. enterica* [23].

*Genomic DNA extraction*

Extraction of DNA was carried out as recommended by the manufacturer of the HiPer® Bacterial Genomic DNA Extraction Teaching Kit (Solution Based), India.

*16S rRNA gene direct sequencing*

Conventional PCR was carried out to analyze all the genes of this study, the confirmatory 16S rRNA gene (621bp) was amplified by using GoTaq® G2 Green Master Mix (Promega, USA). PCR conditions in this study and primer design were used as shown in Tables 1 and 2. Purified PCR products (45µL) from the identified 16S rRNA gene target were forwarded to Macrogen comp. (Korea) for DNA sequencing. Then, using the BioEdit and Mega7 software, consensus sequences were created by aligning the forward and reverse DNA sequences for each sample. In addition, the final sequence from each sample was

**Table 1. PCR conditions of this study**

Gene name		16S rRNA	<i>invA</i>	<i>avrA</i>	<i>sipB</i>	<i>mphA</i>	<i>gyrA</i>
Step conditions							
1 CYCLE	Initial denaturation	95 °C, 5 min					94 °C, 5 min
	Denaturation	94 °C, 40 sec					94 °C, 30 sec
30 CYCLE	Annealing	60 °C	56 °C	58 °C	60 °C	58 °C	58 °C, 30 sec
		40 secs					
	Extension	72 °C, 40 sec					72 °C, 30 sec
1 CYCLE	Final extension	72 °C, 5 min					72 °C, 10 min
—	Holding	4 °C, 10 min					4 °C, 10 min

further analyzed by searching for similar matches in the NCBI gene bank database. This was achieved by employing the BLAST website tool, the BLAST search to assess the similarity.

**Table 2. Primers used in PCR amplification**

Gene name	Primer sequences (5'→3')	Amplicon size (bp)	References
16S rRNA	F primer: GGAAGTACGACACGGTCCAG R primer: CCAGGTAAGGTTCTTCGCGT	671	[23]
	F primer: TTGTTACGGCTATTTGACCA R primer: CTGACTGCTACCTTGCTGATG		
invA	F primer: CCTGTATTGTTGAGCGTCTGG R primer: AGAAGAGCTTCGTTGAATGTCC	425	[25]
	F primer: GGACGCCGCCGGGAAAACTCTC R primer: ACACTCCCGTCGCCCTCACAA		
sipB	F primer: GTGAGGAGGAGCTTCGCGAG R primer: TGCCGAGGACTCGGAGGTC	403	[27]
	F primer: TGGGCAATGACTGGAACA R primer: GGTTGTGCGGCGGGATA		

*Antimicrobial susceptibility test*

The disk diffusion method described by the Clinical and Laboratory Standards Institute (CLSI) [29] was chosen to test antibiotic resistance of *Salmonella* isolates. Ten antimicrobial agents (Bioanalysis, Turkey) were selected from several family groups according to [29], and tested against *Salmonella* isolates: phenicols (chloramphenicol, 30µg), aminoglycosides (gentamicin, 10µg), quinolones (nalidixic acid, 30µg), carbapenems (meropenem and imipenem, 10µg), folate pathway antagonists (trimethoprim-sulfamethoxazole, 30µg), macrolides (azithromycin, 15µg), cephalosporins (ceftazidime and cefotaxime, 30µg), and fosfomycins (fosfomicin, 200µg). *Escherichia coli* ATCC25922 was used as a quality control strain. The isolates were described as susceptible, intermediate, or resistant according to the CLSI [29] guidelines. An isolate was defined as multi-drug resistant (MDR) when showing resistance to three or more different classes of antimicrobials [30].

*Detection of virulence and antimicrobial resistance genes*

The virulence genes *invA* (521bp), *avrA* (425bp), *sipB* (875bp), and the quinolone resistance gene *gyrA* (396bp), macrolide azithromycin resistance gene *mphA* (403bp) were analyzed in isolates that showed resistance to these two antimicrobials only. Bioneer’s master mix (Bioneer’s AccuPower PCR PreMix, Korea) was used for amplification. PCR conditions performed in this study and primer design were used as shown in Table 1 and Table 2.

*Gel electrophoresis*

The agarose powder (1 percent% w/v) was dissolved in 1X TBE buffer. The mixture was microwaved and allowed

to cool at 50 °C before adding 8µl of RedSafe™ (iNtRON/Korea) (0.5 g/ml) to the agarose solution and pouring it onto the tray. After the gel hardened, the comb was removed.


**Results**

*Occurrence of Salmonella spp.*


Among 100 collected samples, 20 isolates (20%) were culture positive for *Salmonella* spp. Within the food sample groups, *Salmonella* spp. was isolated from 16.25% of samples (13 out of 80 food samples), all of which were collected from raw poultry meat and egg groups, whilst the other food groups were *Salmonella* free. Raw frozen chicken breasts had the highest level of contamination with *Salmonella* spp. (60%) followed by raw frozen chicken thighs (40%), raw frozen 9-piece chicken (20%), and eggshell (10%). As regards clinical diarrheal patients, *Salmonella* spp. was isolated from 7 (35%) out of 20 samples. The *S. enterica* isolates from food products identified in this study were *S. enterica* serovar Typhimurium (11 isolates, 84.6%) isolated from raw frozen 9-piece chicken (2 isolates, 15.3%), raw frozen chicken breasts (5 isolates, 38.4%), and raw frozen chicken thighs (4 isolates, 30.7%), *S. enterica* subsp. *enterica* (one isolate from eggs, 7.6%), and *S. enterica* subsp. *diarizonae* (one isolate from raw frozen chicken breasts, 7.6%). *S. enterica* serovar Typhi (4 isolates, 57.1%), and *S. enterica* serovar Typhimurium (3 isolates, 42.8%) were isolated from human diarrheal patients as shown in Table 3. The percentage of each serovar identified in this study is presented in diagrams (Figure 1A and Figure 1B).

**Table 3. Identified isolates and their sources**

Isolates No.	Source of isolates	Type of species
1	Whole raw frozen 9- piece chicken	<i>S. enterica</i> serovar Typhimurium
2		<i>S. enterica</i> serovar Typhimurium
3		<i>S. enterica</i> subsp. <i>diarizonae</i>
4	Raw frozen chicken breasts	<i>S. enterica</i> serovar Typhimurium
5		<i>S. enterica</i> serovar Typhimurium
6		<i>S. enterica</i> serovar Typhimurium
7		<i>S. enterica</i> serovar Typhimurium
8		<i>S. enterica</i> serovar Typhimurium
9	Raw frozen chicken thighs	<i>S. enterica</i> serovar Typhimurium
10		<i>S. enterica</i> serovar Typhimurium
11		<i>S. enterica</i> serovar Typhimurium
12		<i>S. enterica</i> serovar Typhimurium
13	Egg shell	<i>S. enterica</i> subsp. <i>enterica</i>
14	Diarrheal human patients	<i>S. enterica</i> serovar Typhi
15		<i>S. enterica</i> serovar Typhi
16		<i>S. enterica</i> serovar Typhi
17		<i>S. enterica</i> serovar Typhi
18		<i>S. enterica</i> serovar Typhimurium
19		<i>S. enterica</i> serovar Typhimurium
20		<i>S. enterica</i> serovar Typhimurium



**Figure 1A.** Percentage of *S. enterica* serovars identified in raw food samples



**Figure 1B.** Percentage of *S. enterica* serovars identified in clinical human diarrheal samples

As shown in Figures 1A and 1B, the most frequently identified serovars were *S. enterica* serovar Typhimurium (85%) and *S. enterica* serovar Typhi (57%) in raw food and clinical samples, respectively. The confirmatory molecular analysis indicated that all isolates (100%) had the 16S rRNA gene of *Salmonella*.

*Sequencing analysis results*


Several changes in nucleotide sequences were observed in isolates A4, A7, A13, A16, A17, and A19. There was approximately a maximum of 14 differences in base pairs as shown in Figure 2. With the presence of such polymorphisms, differences between isolates belonging to different subspecies and serovars were found.

During the further analysis for bacterial classification and detection of similarity in Figure 2, four clusters of bacterial strains were identified with reference strains. Two of them included AB680591.1 and MZ773245.1, and the other two included isolates A1 and A10. Each of them contained bacterial isolates with the highest similarity and the lowest genetic distance.


Differentiation of the reference strains from those of other serovars and subspecies was carried out as illustrated in Figure 3. It shows that isolates A11, AA2, A1, A8, A20, A12, A5, MZ773245.1, A4, A7, A17, A16, A13, A19, A6, A9, A10, A15, and A18 are more related to strain AB680591.1

**Table 4.** Alignment of partial 16S rRNA gene sequences of *Salmonella* spp. bacteria under consideration with the sequence of the NCBI database

Isolate no.	Result of strain sequencing	Similarity	e-value
A1	<i>S. enterica</i> serovar Typhimurium	100%	0.0
A2		100%	
A3		100%	
A4		99.53%	
A5		100%	
A6		100%	
A7		<i>S. enterica</i> subsp. enterica	
A8	<i>S. enterica</i> serovar Typhimurium	99.50%	
A9	<i>S. enterica</i> serovar Typhi	100%	
A10	<i>S. enterica</i> serovar Typhimurium	99.84%	
A11	<i>S. enterica</i> serovar Typhi	100%	
A12		100%	
A13		99.05%	
A14	<i>S. enterica</i> serovar Typhimurium	100%	
A15		100%	
A16		98.73%	
A17		98.69%	
A18		100%	
A19		100%	
A20	<i>S. enterica</i> subsp. enterica	100%	



**Figure 2.** 16S rRNA gene sequence alignment of *S. enterica* isolates with the related reference sequence of the 16S rRNA gene by BioEdit software. Ref= Reference sequences of the 16S rRNA gene of *S. enterica* strains AB680591.1 S, X80681.1 S, t, and MZ773245.1 S (wild type). The black sign “A” denotes the names of isolates given in Table 4



**Figure 3.** Phylogenetic tree of 16S rRNA gene sequences alignment of *S. enterica* spp. isolates with the related reference sequence of the 16S rRNA gene

and belong to different *S. enterica* serovars while isolates A14 and A13 belonging to *S. enterica* serovar Typhimurium have high similarity to X80681.1 as a common ancestor.

Additionally, the obtained data presented in Table 4 demonstrate the similarity (98–100%) of the isolated bacteria by the 16S rRNA gene, and show that the expected value (e-value) for all *Salmonella* spp. isolates was zero.

*Antimicrobial susceptibility test*

Antimicrobial susceptibility testing of *Salmonella enterica* in this study showed that all isolates tested except one were resistant to one antibiotic at least as shown in Table 5.

In general, the most effective antimicrobials against *Salmonella* isolated from food samples were imipenem (100%), cefotaxime, meropenem and ceftazidime (69.2% each). The highest resistance of all 13 isolates was recorded for nalidixic acid (69.23%), followed by chloramphenicol (53.84%), gentamicin (46.15%), trimethoprim-sulfamethoxazole (46.1%), fosfomycin (46.1%), and azithro-

mycin (38.46%), whilst resistance to both cefotaxime and meropenem was 30.7% as shown in Table 6.

Different *Salmonella* serovars/subspecies tested in this study showed different levels of resistance. It was revealed that 63.63% (7/11) of isolates of *S. enterica* serovar Typhimurium recovered from food were resistant to nalidixic acid and chloramphenicol. Lower resistance was recorded for trimethoprim-sulfamethoxazole and gentamicin, which had the same percentage of 45.45% (6/11), followed by meropenem, cefotaxime, and azithromycin with the same proportion of 36.36% (4/11). Imipenem and ceftazidime were found to be effective antimicrobials against food isolates of *S. enterica* serovar Typhimurium (100% and 76.93%, respectively).

All food isolates (100%) of *S. enterica* subsp. *enterica* (n=1) and *S. enterica* subsp. *diarizonae* (n=1) were susceptible to meropenem, ceftazidime, cefotaxime, and chloramphenicol. Only *S. enterica* subsp. *diarizonae* showed high resistance (100%) to azithromycin, trimethoprim-sulfamethoxazole, gentamicin, nalidixic acid, and fosfo-

**Table 5. Antibiotic resistance of each isolate to ten antibiotics chosen according to CLSI [29]**

sample	Name of serovar/subspecies	NA	SXT	C	CTX	AZM	MEM	IPM	CN	CAZ	FOS	
1	<i>S. enterica</i> serovar Typhimurium	S	S	S	R	S	R	S	R	R	S	MDR
2	<i>S. enterica</i> serovar Typhimurium	S	S	S	R	S	S	S	S	S	S	—
3	<i>S. enterica</i> serovar Typhimurium	S	S	S	S	S	S	S	S	S	S	—
4	<i>S. enterica</i> serovar Typhimurium	S	S	R	S	S	S	S	S	S	R	—
5	<i>S. enterica</i> serovar Typhimurium	R	R	R	R	R	R	S	R	R	R	MDR
6	<i>S. enterica</i> serovar Typhimurium	R	R	S	S	S	R	S	S	S	S	MDR
7	<i>S. enterica</i> subsp. <i>diarizonae</i>	R	R	R	S	R	S	S	R	S	R	MDR
8	<i>S. enterica</i> serovar Typhimurium	R	S	S	S	S	S	S	S	S	S	—
9	<i>S. enterica</i> serovar Typhimurium	R	R	R	S	S	S	S	S	S	R	MDR
10	<i>S. enterica</i> serovar Typhimurium	R	R	R	R	R	R	S	R	R	S	MDR
11	<i>S. enterica</i> serovar Typhimurium	R	S	R	S	R	S	S	R	S	S	MDR
12	<i>S. enterica</i> serovar Typhimurium	R	R	R	S	R	S	S	R	S	R	MDR
13	<i>S. enterica</i> subsp. <i>enterica</i>	R	S	S	S	S	S	S	S	S	S	—
14	<i>S. enterica</i> serovar Typhi	S	R	S	R	S	S	S	R	R	R	MDR
15	<i>S. enterica</i> serovar Typhi	R	S	I	R	R	S	S	S	S	S	MDR
16	<i>S. enterica</i> serovar Typhi	R	S	R	R	S	S	S	S	S	S	MDR
17	<i>S. enterica</i> serovar Typhi	I	R	I	S	R	S	R	R	S	S	MDR
18	<i>S. enterica</i> serovar Typhimurium	R	S	S	R	R	S	S	I	S	R	MDR
19	<i>S. enterica</i> serovar Typhimurium	S	R	R	S	R	R	S	S	R	S	MDR
20	<i>S. enterica</i> serovar Typhimurium	R	S	S	R	R	S	S	I	R	S	MDR

NA: nalidixic acid; SXT: trimethoprim/sulfamethoxazole; C: chloramphenicol, CTX: cefotaxime; AZM: azithromycin; MEM: meropenem; IPM: imipenem; CN: gentamicin; CAZ: ceftazidime; FOS: fosfomycin; S: susceptible; I: intermediate; R: resistant; MDR: multidrug-resistant

mycin. *S. enterica* subsp. *enterica* showed high levels of resistance (100%) to nalidixic acid and high levels of susceptibility (100%) to gentamicin.

Regarding human isolates, the results obtained indicate that in general isolates (*S. enterica* serovar Typhi (n = 4) and *S. enterica* serovar Typhimurium (n = 3)) recovered from the diarrheal patient samples were resistant to azithromycin and cefotaxime showing the same percentage (71.42%; 5/7), while imipenem and meropenem were the ultimate effective antibiotics with the same proportion of susceptible isolates (85.7%; 6/7) as shown in Table7.

As for *Salmonella* serovars, *S. enterica* serovar Typhimurium (n = 3) isolates were completely susceptible (100%) to imipenem, followed by meropenem, fosfomycin, cefotaxime and chloramphenicol with the same proportion (66.67%, 2/3). However, they showed resistance to trimethoprim-sulfamethoxazole, nalidixic acid, and ceftazidime

**Table 6. Percentage of food isolates of *Salmonella* spp. resistant, intermediate and susceptible to antibiotics**

Antibiotics	Food Isolates No.	Resistant (%)	Intermediate (%)	Susceptible (%)
Nalidixic acid (NA)	13	69.2%	0%	30.7%
Trimethoprim/Sulfamethoxazole (SXT)	13	46.1%	0%	53.8%
Chloramphenicol (C)	13	53.8%	0%	46.1%
Cefotaxime (CTX)	13	30.7%	0%	69.2%
Azithromycin (AZM)	13	38.4%	0%	61.5%
Meropenem (MEM)	13	30.7%	0%	69.2%
Imipenem (IPM)	13	0%	0%	100%
Gentamicin (CN)	13	46.1%	0%	53.8%
Ceftazidime (CAZ)	13	30.7%	0%	69.2%
Fosfomycin (FOS)	13	46.1%	0%	53.8%

with the same proportion (66.67%, 2/3). The resistance to gentamicin was intermediate (66.67%, 2/3).

*S. enterica* serovar Typhi (n = 4) isolates exhibited intermediate resistance to chloramphenicol (50%; 2/4). Moreover, they showed resistance to trimethoprim-sulfonamide, nalidixic acid, azithromycin, and gentamicin, all of which had the same ratio (50%; n = 2/4). The highest susceptibility (100%) was observed for meropenem followed by ceftazidime, fosfomycin, and imipenem with the same proportion of 75% (3/4). *S. enterica* serovar Typhi was resistant (75%; 3/4) to cefotaxime.

All isolates except for five isolates from food samples (one from eggshell and four from frozen raw chicken breasts) exhibited multidrug resistance to ≥ 3 antibiotics as shown in Table 5.

This study illustrates a high proportion (77.7%) of food isolates of *S. enterica* serovar Typhimurium (n = 7) consid-

**Table 7. Percentage of clinical *Salmonella* spp. isolates resistant, intermediate and susceptible to antibiotics**

Antibiotics	Clinical Isolates No.	Resistant (%)	Intermediate (%)	Susceptible (%)
Nalidixic acid (NA)	7	57.1%	14.2%	28.5%
Trimethoprim/Sulfamethoxazole (SXT)	7	42.8%	0%	57.1%
Chloramphenicol (CAM)	7	28.5%	28.5%	42.8%
Cefotaxime (CTX)	7	71.4%	0%	28.5%
Azithromycin (AZM)	7	71.4%	0%	28.5%
Meropenem (MEM)	7	14.2%	0%	85.7%
Imipenem (IPM)	7	14.2%	0%	85.7%
Gentamicin (GEN)	7	28.5%	28.5%	42.8%
Ceftazidime (CAZ)	7	42.8%	0%	57.1%
Fosfomycin (FOS)	7	28.5%	0%	71.4%

ered multidrug resistant (MDR). Two isolates were resistant to three antibiotic groups, two isolates were resistant to four antibiotic groups, one isolate was resistant to six antibiotic groups and two isolates were resistant to eight antibiotic groups as shown in Table 8. All (100%) clinical isolates of *S. enterica* serovar Typhimurium (n=3) were MDR: one isolate was resistant to three antibiotic groups, one isolate was resistant to four antibiotic groups and one isolate was resistant to five antibiotic groups as shown in Table 8.

**Table 8. MDR of food and clinical isolates of *S. enterica* serovar Typhimurium**

No. of antimicrobial groups	Food isolates (n=7)		Clinical diarrheal isolates (n=3)	
	Antimicrobial resistance patterns	No. of isolates (%)	Antimicrobial resistance patterns	No. of isolates (%)
Three	Quinolones (NA), Sulfa drug)SXT, Carbapenem (MEM), Cephems (CTX, CAZ), Carbapenem (MEM), Gentamycin (CN).	2 (28.56%)	Quinolones (NA), Cephems (CTX, CAZ), Macrolides (AZM).	1 (14.28%)
Four	Quinolones (NA), Sulfa drug)SXT, Chloramphenicol (C), Fosfomycin (FOS), Quinolones (NA), Chloramphenicol (C), Macrolides (AZM), Gentamycin (CN).	2 (28.56%)	Quinolones (NA), Chloramphenicol (C), Macrolides (AZM), Fosfomycin (FOS).	1 (14.28%)
Five	—	—	Cephems (CAZ), Carbapenem (MEM), Chloramphenicol (C), Macrolides (AZM), Sulfa drug (SXT).	1 (14.28%)
Six	Quinolones (NA), Chloramphenicol (C), Fosfomycin (FOS), Carbapenem (IPM), Macrolides (AZM), Sulfa drug)SXT.	1 (14.28%)	—	—
Eight	Cephems (CAZ, CTX), Carbapenem (MEM), Chloramphenicol (C), Macrolides (AZM), Sulfa drug (SXT), Gentamycin (CN), Quinolones (NA), Fosfomycin (FOS). Cephems (CAZ, CTX), Carbapenem (MEM), Chloramphenicol (C), Macrolides (AZM), Sulfa drug (SXT), Gentamycin (CN), Quinolones (NA), Fosfomycin (FOS).	2 (28.56%)	—	—

*Antibiotic resistance gene detection in S. enterica isolates*

The fluoroquinolone resistance gene (*gyrA*) was associated with all fluoroquinolone (nalidixic acid) resistant isolates (16/16). All nine resistant isolates recovered from food (*S. enterica* serovar Typhimurium, n=7; *S. enterica* subsp. *enterica*, n=1; *S. enterica* subsp. *diarizonae*, n=1) and all seven resistant clinical isolates (*S. enterica* serovar Typhi, n=4; *S. enterica* serovar Typhimurium, n=3) showed the presence of the *gyrA* gene. The azithromycin resistance gene (*mphA*) was associated with about half of the macrolide (azithromycin) resistant isolates (n=6/10). The *mphA* gene was detected in four resistant isolates recovered from food (*S. enterica* serovar Typhimurium, 3/4; and *S. enterica* subsp. *diarizonae*; 1/1) with the proportion equaled 75% and 100%, respectively. Among resistant clinical isolates, 2 out of 5 isolates showed the presence of the *mphA* gene (*S. enterica* serovar Typhi, 1/2; *S. enterica* serovar Typhimurium; 1/3), which accounted for 50% and 33.33%, respectively.

*Virulence gene detection in S. enterica isolates*

The *invA* gene was revealed in 75% (15/20) of all *Salmonella* isolates. Among food isolates, *invA* was found in 9 out of 13 (69.2%) *S. enterica* isolates. Detection of this gene gave negative results in four isolates, including three isolates from raw frozen chicken breasts and one isolate from raw whole 9-piece chicken. All these four negative isolates belonged to *S. enterica* serovar Typhimurium. So, the proportion of *invA* gene detection was 63.6% (7/11) for *S. enterica* serovar Typhimurium isolates, whilst it was 100% for both *S. enterica* subsp. *enterica* (1/1) and *S. enterica* subsp. *diarizonae* (1/1). The *avrA* virulence gene was detected in 90% (18/20) of all *Salmonella* isolates. Among food isolates, *avrA* was detected in 12 out of 13 isolates (92.3%). Only one isolate of *S. enterica* subsp. *enterica* from eggshell showed a negative result of *avrA* gene detection. *S. enterica* serovar Typhimurium and *S. enterica* subsp. *diarizonae* showed a positive result of *avrA* gene detection in all isolates (100%) from local raw poultry meat. The *sipB* virulence gene was found in 95% (19/20) of all *Salmonella* isolates. Regarding local raw poultry meat and eggshell, the *sipB* gene was detected in all (100%) isolates of *S. enterica* serovar Typhimurium, *S. enterica* subsp. *diarizonae*, and *S. enterica* subsp. *enterica*.

All these virulence genes were detected in 85.7% (6/7) of clinical isolates. The proportion of clinical isolates of *S. enterica* serovar Typhi positive for these genes was 25% (1/4).

**Discussion**

To the best of our knowledge, this work is one of the first studies interested in the *Salmonella* prevalence in food and clinical samples in Iraq. The results of *Salmonella* isolates are approximate, with various studies done in many areas in Iraq between 2008 and 2017, with percentage of *Salmonella* isolation ranging from 1.07% to 16%. The great-

est proportion was recorded in Al-Hawijah, while the lowest percentage was recorded in Mosul [31–37].

Different studies on *Salmonella* prevalence were carried out in several countries. In [38] performed in China, *Salmonella* spp. were isolated from 249 out of 664 (37.5%) samples, including 190 (36.7%) chicken, 48 (40.7%) duck and 11 (39.2%) pigeon samples. *Salmonella* prevalence of 13.4% was documented by Rabby [39] in poultry meat retailed in wet- and super-markets in Dhaka city, Bangladesh. Salih *et al.* [40] examined the distribution of *Salmonella* spp. in 121 specimens from diarrheal patients in Duhok, Iraq, and showed that 72 isolates (59.5%) belonged to *Salmonella* spp. Sadeq *et al.* [41] analyzed 40 chicken samples, and found that 14 isolates (35%) belonged to *S. enterica* serovar Typhimurium with the presence of the *invA* gene in 11 (78.5%) out of 14 isolates of *S. enterica* serovar Typhimurium. These results are almost identical to the results of our study since we recorded the presence of seven food isolates of *S. enterica* serovar Typhimurium positive for the *invA* gene with a proportion of 53%. A recent study in Babil, Iraq, was carried out by Obayes *et al.* [42] who collected samples from 120 children with diarrhea and revealed that 58 samples were positive for different *Salmonella* spp. The most common serovar of *Salmonella enterica* in their study was *Salmonella enterica* serovar Typhi (29.3%) and this result agrees with the result of our study, which indicates the presence in the clinical human diarrheal samples of four isolates (57.1%) of *S. enterica* serovar Typhi as the most common serovar.

The most common serovars transferred from animals to humans are *Salmonella* Enteritidis and *Salmonella* Typhimurium. Typhoid fever, paratyphoid fever, food poisoning and gastroenteritis are all disorders caused by *Salmonella* [43]. The 16S rRNA gene sequence is considered an important approach toward identification of bacterial genus, species, and sub-species, and was used in this study as a confirmatory detection test. It is unique for each bacterial organism and can be considered a unique identification gene for bacterial species.

In general, the reasons for *Salmonella* spp. contamination detected in this study are the lack of HACCP control and due diligence. The Iraqi *Central Organization for Standardization and Quality Control* (COSQC, document No. 2270) indicates that the percentage of *Salmonella* growth should be zero in chicken cuts (thighs, breasts, wings), and for this reason the problem of the absence of quality control, HACCP and food handling instructions have to be dissolved and they should be applied in Iraqi slaughterhouses along the poultry processing chain until reaching consumers to prevent an increase in foodborne diseases as much as possible.

Concerning nalidixic acid, other studies also documented *Salmonella* resistance to this antimicrobial agent, and the resistance increased significantly (94.1%) in 2021 compared with the 2018 report (77.3%) indicating more applications of nalidixic acid in both veterinary medicine

and human medicine fields [44, 45]. As for other antibiotics, a study conducted in Bangladesh reported that 95% of the isolates were resistant to azithromycin [46], another study reported low resistance (8%) to chloramphenicol [47]. In [48] 12% of *Salmonella* isolates were resistant to azithromycin and 1% to chloramphenicol, whereas the present study recorded much higher resistance to azithromycin and chloramphenicol compared to previous studies.

In comparison with [49], the partial similarity was noticed, particularly, the susceptibility of the isolates to imipenem. Resistance to SXT was higher in the present study than in [49], which recorded 31.3%, while the study by Velez [50] displayed that *Salmonella* isolates showed higher resistance (100%) to gentamicin than in our study. Moreover, in the current study, nine isolates were resistant to cefotaxime (approximately 30.7%), which is higher than the result reported in China by [14] where 2.44% of *Salmonella* isolates were found to be resistant to cefotaxime.

On other hand, resistance to ceftazidime, meropenem, and fostomycin was in the same line with an Iraqi study [51]. Antibiotic-resistant bacteria can cause life-threatening infections in people and constitute a serious danger to public health and wellbeing. Furthermore, the use of antimicrobials in veterinary medicine may increase the emergence of resistant bacteria harmful to humans and posing a possible threat to public health from zoonotic pathogens, such as *Salmonella*. As a result, the high resistance/MDR in human isolates of *Salmonella* spp. to antibiotics may be caused not by the misuse of drugs and their wrong consumption, but rather this resistance transfers through consumption of foods (e. g., poultry meat) from food-producing animals that received antibiotics on farm to enhance and promote their quality and prevent the growth and transmission of microbes.

Bacteria may acquire resistance genes via mobile genetic elements such as plasmids, which provide the flexibility to a host bacterium and aid in the dissemination and dispersion of these genes among various bacterial populations [52]. In our study, the *gyrA* and *mphA* genes conferring resistance to nalidixic acid and azithromycin, respectively, were detected among isolates that showed the phenotypic resistance. Similarly, there were high percentages of the *gyrA* genes found in nalidixic acid resistant *Salmonella* Albany (92%), *Salmonella* Corvallis (75%), and *Salmonella* Kentucky (85%) isolated in chicken food chains in Cambodia [53].

Quinolones with a broad spectrum of activity have a greater ability to inhibit gyrase in gram-negative bacteria. According to the researchers, antibiotic resistance in *Salmonella* spp. is mostly caused by mutations within the quinolone resistance-determining regions (QRDRs) of the target enzymes DNA gyrase (*gyrA* and *gyrB*) and topoisomerase IV (*parC* and *parE*) [54].

Wang *et al.* [28] identified the *mphA* gene in 15 out of 31 azithromycin-resistant *Salmonella* isolates. This is partially similar to the results of our study. Azithromycin is an

azalide antibacterial drug, which was shown to be equal to chloramphenicol, fluoroquinolones, and extended-spectrum cephalosporins for the treatment of uncomplicated typhoid fever [55]. The *mphA* gene is the key gene involved in *Salmonella* resistance to the macrolide azithromycin. It is typically found on plasmids and spreads rapidly, posing a significant threat to current *Salmonella* infection therapy [56].

*Salmonella* employs several virulence factors expressed at various phases of the disease process to develop a successful infection. A number of these parameters are linked to *Salmonella* Pathogenicity Islands (SPIs) on the *Salmonella* chromosomes [57]. The virulence *invA* gene is involved in *Salmonella* pathogenicity. The *invA* gene acts as a unique biomarker for *Salmonella* identification [58]. Previous studies [59–62] show that the *invA* gene has been found in 100% of *Salmonella* strains, whilst, our study recorded a lower percentage (75%). Likewise, other authors, such as Mthembu *et al.* [21], revealed lower rates (54.4%; 106/195) and Somda *et al.* [63] showed the presence of the *invA* gene in 91% (52/57) of non-typhoidal *Salmonella* isolates from human diarrhea, environment, and lettuce samples in Burkina Faso. Furthermore, Nikiema *et al.* [64] found that the *invA* gene was present in 67% (61/91) of clinical isolates and 60% (9/15) of sandwich samples. In our study, 25% of *Salmonella* isolates did not harbor the *invA* gene and hence would be unable to invade host cells. Salih and Yousif [65] conducted a study in Iraq to detect five virulence factors among four isolates of *S. enterica* serovar Typhimurium isolated previously from three puppies and one adult dog and reported that the *invA* gene was detected in two isolates only. In the same line, another Iraqi study showed that the *invA* gene was detected among eight *Salmonella* strains with a proportion of 50% [66]. Therefore, *Salmonella* might be virulent (*invA*) or avirulent [67]. Furthermore, asymptomatic animals carrying either virulent or avirulent strains might be possible sources of transmission to humans through the food chain, as well as due to their close proximity to people and poor animal effluent management [21, 67]. All strains containing this *invA* gene, which encodes a protein found in bacterial inner membrane and is crucial for invasion of host epithelium, are pathogenic [68,69]. Moreover, a key component of the pathogen's virulence phenotype is the virulence-associated effector protein AvrA of *Salmonella enterica*, which blocks

the first line of defense of the host organism. AvrA expression increases the ability of the bacterium to invade the host [70,71]. Our study indicated that the *avrA* gene was revealed in 90% (18/20) of all *Salmonella* isolates. The other Iraqi researchers Jbar *et al.* [72] detected the *avrA* gene in 100% (30/30) of *Salmonella enterica* isolates. Similarly, an Egyptian study showed that all 6 (100%) *Salmonella* isolates carried the *avrA* gene [69]. This presence of the *avrA* gene in all isolates suggests a higher rate of gastroenteric illnesses in humans that may be transmitted from contaminated food. In addition, Hersh *et al.* [73] established the role of *sipB* in *Salmonella*-induced macrophage death, as well as the possible involvement of caspase-1 in this process. In our study, the *sipB* gene was found in 95% (19/20), while another Iraqi study [42] found that the *sipB* gene occurrence in *Salmonella* isolates was 18.9% (11/120).

Moreover, having regard to the serious roles of these three genes as illustrated briefly above and the outcome of our study, which shows the high occurrence of the *invA*, *avrA*, and *sipB* genes (75%, 90% and 95%, respectively), it is safe to assume that the severity of infections that may occur in the Baghdad population will be increasing.

Differences in the presence of virulence genes in bacterial isolates from this investigation and prior studies might be attributed to geographic circumstances, dietary variables, and the migration of virulence genes through integrons and transposons. In addition to plasmids, [74] indicated that conjugation is a crucial method for the transmission of virulence genes in bacterial groups. These virulence genes, despite their high pathogenicity, are still in circulation, but the variables, as well as the mechanisms of the asymptomatic carriage, are poorly understood. Nonetheless, certain variables, such as a decrease in virulence gene expression or even the expression of bacterial components unique to a carrier, might be blamed. These pathogenicity island-encoded genes are critical in various phases of *Salmonella* pathogenesis [75,76].

## Conclusions

This study shows that *Salmonella enterica* exists in local chicken meat and eggshell with high resistance to antibiotics, including multidrug resistance (75%). In addition, it demonstrates high proportions of *Salmonella enterica* isolates positive for virulence genes (*invA*, *sipB*, *avrA*) responsible for initial invasion of the host cells.

## REFERENCES

1. Cwiek, K., Korzekwa, K., Tabiś, A., Bania, J., Bugla-Płoskońska, G., Wieliczko, A. (2020). Antimicrobial resistance and biofilm formation capacity of *Salmonella enterica* serovar enteritidis strains isolated from poultry and humans in Poland. *Pathogens*, 9(8), Article 643. <https://doi.org/10.3390/pathogens9080643>
2. Pal, M., Ayele, Y. (2020). Emerging role of foodborne viruses in public health. *Biomedical Research International*, 5, 01–04.
3. Kawamoto, S., Bari, M. L. (2015). Emerging and re-emerging foodborne diseases: Threats to human health and global stability. Chapter in a book: *Foodborne Pathogens and Food Safety*. CRC Press, 2015.
4. EFSA (2020). The European Union Summary Report on Antimicrobial Resistance in zoonotic and indicator bacteria from humans, animals and food in 2017/2018. *EFSA*, 18(3), Article 6007. <https://doi.org/10.2903/j.efsa.2020.6007>
5. Khalaf, Z. Z. (2018). Comparative study of antibiofilm activity of lime juice and lithium dioxide nanoparticles against *Salmonella* isolated from local cheese. *Journal of Biotechnology Research Center*, 12(2), 82–94. <https://doi.org/10.24126/jobrc.2018.12.2.542>
6. ECDC. (2020). *Salmonella* the Most Common Cause of Foodborne Outbreaks in the European Union. Retrieved from

- <https://www.ecdc.europa.eu/en/news-events/Salmonella-most-common-cause-foodborne-outbreaks-european-union> Accessed December 15, 2022
7. Jajere, S. M. (2019). A review of *Salmonella enterica* with particular focus on the pathogenicity and virulence factors, host specificity and antimicrobial resistance including multi-drug resistance. *Veterinary World*, 12(4), 504–521. <https://doi.org/10.14202/vetworld.2019.504-521>
  8. Muller, K., Aabo, S., Birk, T., Mordhorst, H., Bjarnadottir, B., Agersø, Y. (2012). Survival and growth of epidemically successful and unsuccessful *Salmonella enterica* clones after freezing and dehydration. *Journal of Food Protection*, 75(3), 456–464. <https://doi.org/10.4315/0362-028X.JFP-11-167>
  9. Rosenkrantz, J. T., Aarts, H., Abee, T., Rolfe, M. D., Knudsen, G. M., Nielsen, M. B. et al. (2013). Non-essential genes form the hubs of genome scale protein function and environmental gene expression networks in *Salmonella enterica* serovar Typhimurium. *BMC Microbiology*, 13(1), Article 294. <https://doi.org/10.1186/1471-2180-13-294>
  10. Brooks, G. F., Butel, J. S., Morse, S. A. (2004). Cultivation of Microorganisms. Chapter in a book: Medical Microbiology. New York, Cengage Publisher Services, 2004.
  11. Tindall, B. J., Grimont, P. A. D., Garrity, G. M., Euzéby, J. P. (2005). Nomenclature and taxonomy of the genus *Salmonella*. *International Journal of Systematic and Evolutionary Microbiology*, 55(1), 521–524. <https://doi.org/10.1099/ijs.0.63580-0>
  12. Brenner, F. W., Villar, R. G., Angulo, F. J., Tauxe, R., and Swaminathan, B. (2000). *Salmonella* Nomenclature. *Journal of Clinical Microbiology*, 38(7), 2465–2467. <https://doi.org/10.1128/JCM.38.7.2465-2467.2000>
  13. Harb, A., Habib, I., Mezal, E. H., Kareem, H. S., Laird, T., O’Dea, M. et al. (2018). Occurrence, antimicrobial resistance and whole-genome sequencing analysis of *Salmonella* isolates from chicken carcasses imported into Iraq from four different countries. *International Journal of Food Microbiology*, 284, 84–90. <https://doi.org/10.1016/j.ijfoodmicro.2018.07.007>
  14. Ma, Y., Xu, X., Gao, Y., Zhan, Z., Xu, C., Qu, X. et al. (2020). Antimicrobial resistance and molecular characterization of *Salmonella enterica* serovar Corvallis isolated from human patients and animal source foods in China. *International Journal of Food Microbiology*, 335, Article 108859. <https://doi.org/10.1016/j.ijfoodmicro.2020.108859>
  15. Lai, J., Wu, C., Wu, C., Qi, J., Wang, Y., Wang, H. et al. (2014). Serotype distribution and antibiotic resistance of *Salmonella* in food-producing animals in Shandong province of China, 2009 and 2012. *International Journal of Food Microbiology*, 180, 30–38. <https://doi.org/10.1016/j.ijfoodmicro.2014.03.030>
  16. Yang, B., Qiao, L., Zhang, X., Cui, Y., Xia, X., Cui, S. et al. (2013). Serotyping, antimicrobial susceptibility, pulse field gel electrophoresis analysis of *Salmonella* isolates from retail foods in Henan Province, China. *Food Control*, 32(1), 228–235. <https://doi.org/10.1016/j.foodcont.2012.11.022>
  17. Zhu, Y., Lai, H., Zou, L., Yin, S., Wang, C., Han, X. et al. (2017). Antimicrobial resistance and resistance genes in *Salmonella* strains isolated from broiler chickens along the slaughtering process in China. *International Journal of Food Microbiology*, 259, 43–51. <https://doi.org/10.1016/j.ijfoodmicro.2017.07.023>
  18. Harb, A., O’Dea, M., Hanan, Z. K., Abraham, S., Habib, I. (2017). Prevalence, risk factors and antimicrobial resistance of *Salmonella* diarrhoeal infection among children in Thi-Qar Governorate, Iraq. *Epidemiology and Infection*, 145(16), 3486–3496. <https://doi.org/10.1017/S0950268817002400>
  19. Fabrega, A., Vila, J. (2013). *Salmonella enterica* serovar typhimurium skills to succeed in the host: Virulence and regulation. *Clinical Microbiology Reviews*, 26(2), 308–341. <https://doi.org/10.1128/cmr.00066-12>
  20. Lou, L., Zhang, P., Piao, R., Wang, Y. (2019). *Salmonella* Pathogenicity Island 1 (SPI-1) and its complex regulatory network. *Frontiers in Cellular and Infection Microbiology*, 9, Article 270. <https://doi.org/10.3389/fcimb.2019.00270>
  21. Mthembu, T. P., Zishiri, O. T., El Zowalaty, M. E. (2019). Detection and molecular identification of *Salmonella* virulence genes in livestock production systems in South Africa. *Pathogens*, 8(3), Article 124. <https://doi.org/10.3390/pathogens8030124>
  22. Mostafa, L., Aioub, K., Mousa, M., Ahmed, A. (2019). Effect of some essential oils on *Salmonella* Kentucky isolated from quail meat retailed in Alexandria markets. *Alexandria Journal of Veterinary Sciences*, 61(1), Article 179. <https://doi.org/10.5455/ajvs.29683>
  23. Kaabi, H. K. J. A., AL-Yassari, A. K. S. (2019). 16SrRNA sequencing as tool for identification of *Salmonella* spp. isolated from human diarrhea cases. *Journal of Physics: Conference Series*, 1294(6), Article 062041. <https://doi.org/10.1088/1742-6596/1294/6/062041>
  24. Van der Velden, A. W., Lindgren, S. W., Worley, M. J., Heffron, F. (2000). *Salmonella* pathogenicity island 1-independent induction of apoptosis in infected macrophages by *Salmonella enterica* serotype Typhimurium. *Infection and Immunity*, 68(10), 5702–5709. <https://doi.org/10.1128/iai.68.10.5702-5709.2000>
  25. Huehn, S., La Ragione, R. M., Anjum, M., Saunders, M., Woodward, M. J., Bunge, C. et al. (2010). Virulotyping and antimicrobial resistance typing of *Salmonella enterica* serovars relevant to human health in Europe. *Foodborne Pathogens and Disease*, 7(5), 523–535. <https://doi.org/10.1089/fpd.2009.0447>
  26. Skyberg, J. A., Logue, C. M., Nolan, L. K. (2009). Virulence genotyping of *Salmonella* spp. with multiplex PCR. *Avian Diseases*, 50(1), 77–81. <https://doi.org/10.1637/7417.1>
  27. Gomes, C., Ruiz-Roldán, L., Mateu, J., Ochoa, T. J., Ruiz, J. (2019). Azithromycin resistance levels and mechanisms in *Escherichia coli*. *Scientific Reports*, 9(1), Article 6089. <https://doi.org/10.1038/s41598-019-42423-3>
  28. Wang, J., Li, Y., Xu, X., Liang, B., Wu, F., Yang, X. et al. (2017). Antimicrobial resistance of *Salmonella enterica* serovar Typhimurium in Shanghai, China. *Frontiers in Microbiology*, 8, Article 510. <https://doi.org/10.3389/fmicb.2017.00510>
  29. Clinical and Laboratory Standards Institute (CLSI). 2021. Performance standards for antimicrobial susceptibility testing. 31th (ed.). CLSI document M100. Wayne, USA.
  30. Abraham, S., Groves, M. D., Trott, D. J., Chapman, T. A., Turner, B., Hornitzky, M. et al. (2014). *Salmonella enterica* isolated from infections in Australian livestock remain susceptible to critical antimicrobials. *International Journal of Antimicrobial Agents*, 43(2), 126–130. <https://doi.org/10.1016/j.ijantimicag.2013.10.014>
  31. Alrifai, S. B., Alsaadi, A., Mahmood, Y. A., Ali, A. A. (2009). Prevalence and etiology of nosocomial diarrhoea in children < 5 years in Tikrit teaching hospital. *EMHJ – Eastern Mediterranean Health Journal*, 15(5), 1111–1118. <https://doi.org/10.26719/2009.15.5.111>
  32. Rabatti, A.A., Rasheed, N.E. (2009). Etiology of bloody diarrhea among children admitted to maternity and children’s Hospital-Erbil. *Al-Kindy College Medical Journal*, 4(2), 19–24.
  33. Al-Mosawi, G.J., Al-Haris, F.M. (2008). The aetiology of bloody diarrhea in children of najaf governorate. *Kufa Medical Journal*, 11(1), 224–233.
  34. Al-Taie, H. H. I. (2009). Prevalence rate of *Salmonella* in Babylon province. *The Iraqi Journal of Veterinary Medicine*, 33(2), 152–157. <https://doi.org/10.30539/iraqijvm.v33i2.705>
  35. Al-Juboory, Y.H., Zenad, M.M., Hassen, R. H. (2014). Prevalence of *Salmonella* Serotypes in Diarrheic and Non-Diarrheic Patients in Mosul-Iraq. *Kerbala Journal of Medicine*, 7(2), 1937–1944.
  36. Saleh, M.B., Hanan, Z.K., Mezal, E.H. (2016). Antimicrobial resistance, virulence profiles of *Salmonella enterica* serovar Typhimurium isolated from diarrheal children in Thi-Qar province during 2015. *University of Thi-Qar Journal of Science*, 6(1), 3–8.
  37. Abbas, K., Al-Wattar, W., Hasan, S., Kasim, H., Jasim, A. (2017). The incidence of Shigella and *Salmonella* in the stool of pediatric patients. *Iraqi Journal of Public Health*, 1(3), 2521–7267. <https://doi.org/10.22317/ijph.12201705>
  38. Yang, X., Huang, J., Zhang, Y., Liu, S., Chen, L., Xiao, C. et al. (2020). Prevalence, abundance, serovars and antimicrobial resistance of *Salmonella* isolated from retail raw poultry meat in China. *Science of The Total Environment*, 713, Article 136385. <https://doi.org/10.1016/j.scitotenv.2019.136385>
  39. Rabby, M. R. I., Shah, S. T., Miah, M. I., Islam, M. S., Khan, M. A. S., Rahman, M. S. et al. (2021). Comparative analysis of bacteriological hazards and prevalence of *Salmonella* in poultry-meat retailed in wet- and super-markets in Dhaka city, Bangladesh. *Journal of Agriculture and Food Research*, 6, Article 100224. <https://doi.org/10.1016/j.jafr.2021.100224>
  40. Salih, D., Abdo, J., Saadi, A. (2019). Molecular identification of *Salmonella enterica* from patients with diarrhea in Duhok governorate Kurdistan region/Iraq. *Journal of Duhok University*, 22(1), 148–154. <https://doi.org/10.26682/sjuod.2019.22.1.16>
  41. Sadeq, J. N., Esmaeel, J. R., Neama, A. A. (2017). Molecular detection of *invA*, *ssaP* in *Salmonella typhimurium* isolated from chicken in Al-Qadisiyah Province. *AL-Qadisiyah Journal of Veterinary Medicine Sciences*, 16(2), 8–13. <https://doi.org/10.29079/vol16iss2art436>
  42. Obayes, M. S., Al-Bermani, O. K., Rahim, S. A. (2020). Genetic detection of *invA*, *sipB*, *SopB* and *sseC* genes in *Salmonella* spp. isolated from diarrheic children patients. *Eurasian Journal of Biosciences*, 14(2), 3085–3091.

43. Preethi, B., Shanthi, V., Ramanathan, K. (2015). Investigation of nalidixic acid resistance mechanism in *Salmonella enterica* using molecular simulation techniques. *Applied Biochemistry and Biotechnology*, 177, 528–540. <https://doi.org/10.1007/s12010-015-1760-6>
44. Fardsanei, F., Soltan Dallal, M. M., Douraghi, M., Memariani, H., Bakhshi, B., Zahraei Salehi, T. Z. et al. (2018). Antimicrobial resistance, virulence genes and genetic relatedness of *Salmonella enterica* serotype Enteritidis isolates recovered from human gastroenteritis in Tehran, Iran. *Journal of Global Antimicrobial Resistance*, 12, 220–226. <https://doi.org/10.1016/j.jgar.2017.10.005>
45. Wei, B., Shang, K., Cha, S.-Y., Zhang, J.-F., Jang, H.-K., Kang, M. (2021). Clonal dissemination of *Salmonella enterica* serovar albany with concurrent resistance to ampicillin, chloramphenicol, streptomycin, sulfisoxazole, tetracycline, and nalidixic acid in broiler chicken in Korea. *Poultry Science*, 100(7), Article 101141. <https://doi.org/10.1016/j.psj.2021.101141>
46. Ahsan, S., Rahman, S. (2019). Azithromycin resistance in clinical isolates of *Salmonella enterica* serovars typhi and paratyphi in Bangladesh. *Microbial Drug Resistance*, 25(1), 8–13. <https://doi.org/10.1089/mdr.2018.0109>
47. Mthembu, T. P., Zishiri, O. T., El Zowalaty, M. E. (2019). Molecular detection of multidrug-resistant *Salmonella* isolated from livestock production systems in South Africa. *Infection and Drug Resistance*, 12, 3537–3548. <https://doi.org/10.2147/idr.s211618>
48. Girish, R., Kumar, A., Khan, S., Dinesh, K. R., Karim, S. (2013). Revised ciprofloxacin breakpoints for *Salmonella*: Is it time to write an obituary? *Journal of Clinical and Diagnostic Research*, 7(11), 2467–2469. <https://doi.org/10.7860/JCDR/2013/7312.3581>
49. Yu, X., Zhu, H., Bo, Y., Li, Y., Zhang, Y., Liu, Y. (2021). Prevalence and antimicrobial resistance of *Salmonella enterica* subspecies enterica serovar Enteritidis isolated from broiler chickens in Shandong Province, China, 2013–2018. *Poultry Science*, 100(2), 1016–1023. <https://doi.org/10.1016/j.psj.2020.09.079>
50. Velez, D. C., Rodríguez, V., & García, N. V. (2017). Phenotypic and genotypic antibiotic resistance of *Salmonella* from chicken carcasses marketed at Ibagué, Colombia. *Brazilian Journal of Poultry Science*, 19, 347–354. <http://doi.org/10.1590/1806-9061-2016-0405>
51. Hasan, T. O. (2021). Genomic diversity analysis of *Salmonella* spp. from broiler and layer flocks and their feed and water in Karbala, Iraq. Ph.D. thesis, University of Baghdad, 2021. <https://doi.org/10.13140/RG.2.2.23584.71682>
52. Blair, J. M. A., Webber, M. A., Baylay, A. J., Ogbolu, D. O., Piddock, L. J. V. (2014). Molecular mechanisms of antibiotic resistance. *Nature Reviews Microbiology*, 13(1), 42–51. <https://doi.org/10.1038/nrmicro3380>
53. Vuthy, Y., Lay, K. S., Seiha, H., Kerleguer, A., Aidara-Kane, A. (2017). Antibiotic susceptibility and molecular characterization of resistance genes among *Escherichia coli* and among *Salmonella* subsp. in chicken food chains. *Asian Pacific Journal of Tropical Biomedicine*, 7(7), 670–674. <https://doi.org/10.1016/j.apjtb.2017.07.002>
54. Preethi, B., Shanthi, V., Ramanathan, K. (2015). Investigation of nalidixic acid resistance mechanism in *Salmonella enterica* using molecular simulation techniques. *Applied Biochemistry and Biotechnology*, 177(2), 528–540. <https://doi.org/10.1007/s12010-015-1760-6>
55. Sharma, P., Kumari, B., Dahiya, S., Kulsum, U., Kumar, S., Manral, N. et al. (2019). Azithromycin resistance mechanisms in typhoidal *Salmonellae* in India: A 25 years analysis. *Indian Journal of Medical Research*, 149(3), 404–411. [https://doi.org/10.4103/ijmr.IJMR\\_1302\\_17](https://doi.org/10.4103/ijmr.IJMR_1302_17)
56. Wang, H., Cheng, H., Huang, B., Hu, X., Chen, Y., Zheng, L., et al. (2023). Characterization of resistance genes and plasmids from sick children caused by *Salmonella enterica* resistance to azithromycin in Shenzhen, China. *Frontiers in Cellular and Infection Microbiology*, 13, 343. <https://doi.org/10.3389/fcimb.2023.11161722>
57. Golubeva, Y. (2010). Regulation of virulence in *Salmonella enterica*. Ph.D. thesis, University of Illinois at Urbana-Champaign 2010.
58. Li, Q., Cheng, W., Zhang, D., Yu, T., Yin, Y., Ju, H. et al. (2012). Rapid and sensitive strategy for *Salmonella* detection using an *InvA* gene-based electrochemical DNA sensor. *International Journal of Electrochemical Science*, 7(1), 844–856.
59. Chaudhary, J.H., Nayak, J.B., Brahmabhatt, M.N., Makwana, P.P. (2015). Virulence genes detection of *Salmonella* serovars isolated from pork and slaughterhouse environment in Ahmedabad, Gujarat. *Veterinary World*, 8(1), 121–124. <https://doi.org/10.14202/vetworld.2015.121-124>
60. Gassama-Sow, A., Wane, A.A., Canu, N.A., Uzzau, S., Aidara-Kane, A., Rubino, S. (2006). Characterization of virulence factors in the newly described *Salmonella enterica* serotype Keurmassar emerging in Senegal (sub-Saharan Africa). *Epidemiology and Infection*, 134(4), 741–743. <https://doi.org/10.1017/S0950268805005807>
61. Borges, K.A., Furian, T.Q., Borsoi, A., Moraes, H.L., Salle, C.T., Nascimento, V.P. (2013). Detection of virulence-associated genes in *Salmonella* Enteritidis isolates from chicken in South of Brazil. *Pesquisa Veterinária Brasileira*, 33, 1416–1422.
62. Deguenon, E., Dougnon, V., Lozes, E., Maman, N., Agbankpe, J., Abdel-Massih, R.M. et al. (2019). Resistance and virulence determinants of faecal *Salmonella* spp. isolated from slaughter animals in Benin. *BMC Research Notes*, 12(1), Article 317. <https://doi.org/10.1186/s13104-019-4341-x>
63. Somda, N.S., Bonkougou, I.J.O., Sambe-Ba, B., Drabo, M.S., Wane, A.A., Sawadogo-Lingani, H. et al. (2021). Diversity and antimicrobial drug resistance of non-typhoid *Salmonella* serotypes isolated in lettuce, irrigation water and clinical samples in Burkina Faso. *Journal of Agriculture and Food Research*, 5, Article 100167. <https://doi.org/10.1016/j.jafr.2021.100167>
64. Nikiema, M.E.M., Kakou-Ngazon, S., Ky/Ba, A., Sylla, A., Bako, E., Addablah, A.Y.A. et al. (2021). Characterization of virulence factors of *Salmonella* isolated from human stools and street food in urban areas of Burkina Faso. *BMC Microbiology*, 21(1), Article 338. <https://doi.org/10.1186/s12866-021-02398-6>
65. Salih, W., Yousif, A.A. (2018). Molecular detection of *Salmonella typhimurium* isolated from canine feces by PCR. *Advances in Animal and Veterinary Sciences*, 6(12), 542–547. <https://doi.org/10.17582/journal.aavs/2018/6.12.542.547>
66. Kadry, M., Nader, S.M., Dorgham, S.M., Kandil, M.M. (2019). Molecular diversity of the *invA* gene obtained from human and egg samples. *Veterinary World*, 12(7), Article 1033. <https://doi.org/10.14202/vetworld.2019.1033-1038>
67. Ahmer, B. M. M., Gunn, J. S. (2011). Interaction of *Salmonella* spp. with the intestinal microbiota. *Frontiers in Microbiology*, 2, Article 101. <https://doi.org/10.3389/fmicb.2011.00101>
68. Hassan, K. I., Saleh, S. H. (2016). A rapid method for PCR based on detection of *Salmonella* spp. and *Staphylococcus aureus* in spiked and naturally contaminated food. *Iraqi Journal of Agricultural Sciences*, 47(7 – special issue), 66–73.
69. Oludairo, O.O., Kwaga, J.K.P., Dzikwi, A. A., Junaid, K. (2013). Detection of *invA* virulence gene by polymerase chain reaction (PCR) in *Salmonella* spp. isolated from captive wildlife. *Bio-Genetics Journal*, 1(1), 12–14.
70. Lu, R., Wu, S., Liu, X., Xia, Y., Zhang, Y. -G., Sun, J. (2010). Chronic effects of a *Salmonella* type III secretion effector protein *AvrA* in vivo. *PLoS ONE*, 5(5), Article e10505. <https://doi.org/10.1371/journal.pone.0010505>
71. Saber, A. S., Abeer, H.A., (2019). Molecular characterization of *Salmonella* species isolated from chicken table egg content. *Assiut Veterinary Medical Journal*, 65(162), 83–92. <https://doi.org/10.21608/avmj.2019.168950>
72. Jbar, H.S., Al-Janabi, H. S. Kadhim, M.J. (2021). Molecular detection of some outer membrane proteins related to immune resistance in *Salmonella enterica*. *Turkish Journal of Physiotherapy and Rehabilitation*, 32(3), 18739–18743.
73. Hersh, D., Monack, D. M., Smith, M. R., Ghorri, N., Falkow, S., Zychlinsky, A. (1999). The *Salmonella* invasin *SipB* induces macrophage apoptosis by binding to caspase-1. *Proceedings of the National Academy of Sciences*, 96(5), 2396–2401.
74. Cabezon, E., Ripoll-Rozada, J., Pena, A., de la Cruz, F., Arechaga, I. (2015). Towards an integrated model of bacterial conjugation. *Microbiol. Rev. FEMS Microbiology Reviews*, 39(1), 81–95. <https://doi.org/10.1111/1574-6976.12085>
75. Ahmet, N. A., Bissoume, S. B., Abdoulaye, S., Amadou, D., Khota, F. N., Aziz, W. A. et al. (2020). Phenotypic and genotypic characterization of *Salmonella* isolated from Asymptomatic Carriers in the Suburb of Dakar. *Journal of Tropical Diseases and Public Health*, 8, Article 350. <https://doi.org/10.35248/2329-891X.20.8.346>
76. Allaith, S. A., Abdel-aziz, M. E., Thabit, Z. A., Altemimi, A. B., Abd El-Ghany, K., Giuffrè, A. M. et al. (2022). Screening and molecular identification of lactic acid bacteria producing  $\beta$ -glucan in Boza and Cider. *Fermentation*, 8(8), Article 350. <https://doi.org/10.3390/fermentation8080350>

#### AUTHOR INFORMATION

**Zahraa A. AlShaheeb**, MSc, Assistant lecturer, Laboratory Scientist, Iraqi Ministry of Health. Baghdad 64074, Iraq. Tel.: +9647830602282, E-mail: zahraa.abdulkareem@ced.nahrainuniv.edu.iq  
ORCID: <https://orcid.org/0000-0002-0193-9218>

**Zaid A. Thabit**, PhD, Assistant professor, Academic Staff, Environmental Biotech Department, Biotechnology Research Center, Al-Nahrain University. Baghdad 64074, Iraq.

Tel.: +9647700326685, E-mail: zaid.alrawi@brc.nahrainuniv.edu.iq

ORCID: <https://orcid.org/0000-0002-3431-7562>

\* corresponding author

**Asma G. Oraibi**, PhD, Professor, Academic Staff, Plant Biotechnology Department, College of Biotechnology, Al-Nahrain University, Baghdad 64074, Iraq.

Tel.: +9647819905976, E-mail: asma.ghatea@nahrainuniv.edu.iq

ORCID: <https://orcid.org/0000-0002-0533-1349>

**Ahmed A. Baioumy**, PhD, Assistant Professor, Food Science Department, Faculty of Agriculture, Cairo University. 12613, 1 Gamaa Street, Giza, Egypt. Tel.: +2-0122-838-44-78, E-mail: ahmedadel35@agr.cu.edu.eg

ORCID: <https://orcid.org/0000-0002-2909-603X>

**Tarek G. Abdelmaksoud**, PhD, Assistant Professor, Food Science Department, Faculty of Agriculture, Cairo University. 12613, 1 Gamaa Street, Giza, Egypt. Tel.: +2-0110-144-12-80, E-mail: tareekgamal\_88@agr.cu.edu.eg

ORCID: <https://orcid.org/0000-0002-7012-6667>

All authors bear responsibility for the work and presented data.

All authors made an equal contribution to the work.

The authors were equally involved in writing the manuscript and bear the equal responsibility for plagiarism.

The authors declare no conflict of interest.